

Use of progestagens during early pregnancy

G. DANTE, V. VACCARO, F. FACCHINETTI

Mother-Infant Department, Unit of Obstetrics, University of Modena and Reggio Emilia, Italy.

Correspondence at: fabio.facchinetti@unimore.it

Abstract

The term “progestagens” covers a group of molecules including both the natural female sex hormones Progesterone and 17-hydroxy Progesterone as well as several synthetic forms, all displaying the ability to bind Progesterone receptors.

Several studies have used Progesterone and related steroids in the attempt to prevent spontaneous miscarriage, and treat recurrent miscarriage.

The present paper aims to provide a comprehensive review of the literature on progestagens effects during early pregnancy. We looked only at the results from randomized controlled trials. We found and analyzed 15 trials on the prevention of recurrent miscarriage and 2 trials on the treatment of miscarriage. The results demonstrated that there is no evidence to support the routine use of progestagens for the treatment of threatened miscarriage.

Key words: Miscarriage, progestagens, Progesterone, 17 α OH Progesterone, 17 α OH Progesterone Caproate, recurrent miscarriage.

Introduction

Progesterone (Fig. 1) is a steroid hormone which plays a crucial role in each step of human pregnancy. In early pregnancy Progesterone is produced by the corpus luteum whose duration has been estimated 12 ± 2 days. This organ is fundamental for pregnancy maintenance until the placenta (syncytiotrophoblast) takes over its function at 7-9th week of gestation, just after the expression of major histocompatibility complex antigens is suppressed in extra-embryonic fetal tissue.

Progesterone is an essential hormone in the process of reproduction. Indeed, it induces secretory changes in the lining of the uterus and is essential for a successful implantation of the embryo. Moreover, Progesterone modulates the immune response of the mother to prevent rejection of the embryo, and enhances uterine quiescence and suppresses uterine contractions. Therefore it is theoretically plausible that P supplementation may reduce the risk of miscarriage in women with a history of recurrent miscarriages. Several studies have used Progesterone and related steroids (progestagens- Fig. 1) in the attempt to prevent spontaneous miscarriage and

to increase the embryo implantation rates in assisted reproduction programmes. The term “progestagens” cover a group of molecules including both the natural female sex hormones Progesterone and 17-hydroxy Progesterone (17P) as well as several synthetic forms, all displaying the ability to bind Progesterone receptors.

Although pharmacokinetics and pharmacodynamics features of progestagens have been studied, their use in human pregnancy remains controversial, i.e. the way of administration. Indeed, progestagens could be administered by three routes: orally, vaginally or intramuscularly. Oral administration guarantees optimal compliance by patients but shows many disadvantages; this route also results in side effects such as nausea, headache and sleepiness. The vaginal route results in higher concentrations in the uterus but does not reach high and constant blood levels. The drug administered intramuscularly occasionally induces non-septic abscesses, although it is the only route which results in optimal blood levels (Whitehead et al., 1990; Szabo and Szilagyi, 1996; Cunningham, 2001; Di Renzo et al., 2005; Christian et al., 2007). Several reports hypothesized an association between intrauterine exposure to

Progesterone




17 α OH Progesterone17 α OH Progesterone Caproate

Fig. 1. — Progesterone and Progestagens

progestagens in the first trimester of pregnancy and genital abnormalities in male and female fetuses. This was due to the possible up-regulation of androgen receptor operated by pharmacological doses of these steroids. However, maternal safety of progestagens has been reported in different trials. Neonatal safety has been evaluated in only one trial where mothers have been treated with 17 Progesterone. No effects of general health status, external genitalia, and psychomotor development have been reported at follow-up. Since the paucity of data, ongoing trials are encouraged to include the follow-up of neonates in their study design (Mosby, 2001).

The present paper aims to provide a comprehensive view of the literature on the effects of progestagens during early pregnancy. We describe the effects of progestagens for preventing recurrent miscarriages and managing threatened miscarriage.

Miscarriage is defined as pregnancy loss before 23 weeks' gestation, based on the first day of the last menstrual period. Miscarriage is associated with considerable physical and psychological morbidity, particularly in developing countries (World Health Organization, 1992). Haemorrhage into the decidua basalis and necrotic changes in the tissues adjacent to the bleeding usually accompany abortion; the ovum becomes detached and stimulates uterine contractions that result in expulsion (Cunningham, 2001).

Recurrent miscarriage has been defined as 3 or more consecutive episodes of spontaneous pregnancy losses with the same biological father (World Health Organization, 1992). Extensive investigation of women involved will fail to find a recognisable cause in up to half of the cases. Luteal phase defects, immunotolerance derangements, chromosomal anomalies and endocrine disorders are the most common recognisable causes. Accepting an independent risk of miscarriage to be 15%, a second loss could be calculated to occur at a rate of 2-3% while a third loss is expected in 0.34% of women.

More than 80% of miscarriages occur before the 12th week, and the rate decreases rapidly thereafter. It is a common complication of pregnancy occurring in 15% to 20% of all clinically recognized pregnancies with 1% to 2% of couples suffering a recurrent early loss. It is thought, however, that the true incidence of early spontaneous miscarriage may be much higher. Indeed, after implementation of assisted reproduction techniques we are able to detect biochemical pregnancies and therefore to conclude that the rate of early spontaneous miscarriage in the first trimester is higher than expected (Everett, 1997).

Chromosomal anomalies cause at least half of these early miscarriages. The risk increases with parity and at the extremes of maternal age (>34 years and <20 years old). Other risk factors for miscarriage include maternal infections (such as vaginitis, urinary tract infections), endocrine abnormalities (diabetes mellitus, hypothyroidism, pituitary adenoma), insufficient production of Progesterone by the corpus luteum (short and/or abnormal luteal phase), polycystic ovary syndrome, drug use, environmental factors, maternal autoimmune disorders (such as anti-phospholipids antibodies), previous history of two or more miscarriages and acquired or inherited uterine malformations (Szabo and Szilagyi, 1996). Despite all the above recognised or supposed factors, the cause of miscarriage cannot be identified in almost 50% of cases. Threatened miscarriage manifests itself through vaginal bleeding, with or without abdominal pain, while the cervix is closed and the fetus is viable inside the uterine cavity. The introduction of ultrasound scans in the management of bleeding in early pregnancy has improved the diagnosis tremendously (Hemminki, 1998).

During the past 50 years several trials investigated the use of progestagens for the prevention of miscarriage. Actually the therapeutic value of progestagens remains to be established. This might be due to the poor design of the studies which evaluated hormone

effectiveness. Moreover, a lot of different aetiologies are associated with threatened miscarriages and heterogeneity of studies has not been accounted for.

Methods

Literature searches were performed in The Cochrane Library, only randomized controlled trials (RCTs) were considered. The search terms were as follows: “progestagens”, “miscarriage”, “recurrent miscarriage”. We performed a search about all types of progestagens, their applications and potential effects for preventing recurrent miscarriage and for managing miscarriage, regardless of the dose, duration or route of administration compared with placebo, no treatment or other intervention.

All sources of information were read and evaluated by one of the authors (GD), and were later independently checked by another author (VV).

Results

Two recent Cochrane Reviews (Haas and Ramsey, 2009; Wahabi et al., 2011) selected the randomized controlled trials (RCT) or quasi-randomized con-

trolled trials published between 1966 and 2006. They included all types of progestagens in the prevention of miscarriage (Haas and Ramsey, 2009) and for the treatment of threatened miscarriage (Wahabi et al., 2011), regardless of the dose, duration or route of administration and compared with placebo, no treatment or other interventions. No further RCTs were published after those included in the Cochrane Review.

Wahabi et al. (2011) selected the trials published in 7 databases and references from relevant articles. The authors included only the trials where the administration of progestagens started before pregnancy and continued during pregnancy. Twenty-nine studies were potentially eligible for inclusion, of these, 15 studies were included after applying the inclusion criteria. The characteristics of these RCTs are described in Table 1.

Haas and Ramsey (2009) selected the RCTs or quasi-randomized controlled trials that appeared in 30 journals and the proceedings of major conferences describing all types of progestagens in the treatment of threatened miscarriage. Participants were all pregnant women, with a history of threatened miscarriage and a confirmed viable pregnancy.

Table 1. — Characteristics of randomized controlled trials analyzed for preventing recurrent miscarriage (RCT: Randomized Controlled Trial, PTB: preterm birth).

Authors/year	Population studied	Intervention	Comparison	Outcome measured
El Zibdeh et al. 2005	Pregnant women (aged <35 years) with at least 3 consecutive unexplained abortions with same partner N = 180/180	- Dihydrogesterone - Oral - Until miscarriage or 12 weeks - 10 mg twice daily	RCT vs no treatment	Miscarriage, PTB, fetal malformations
Goldzieher et al. 1964	Women who never had a term pregnancy and with a history of 2 or more miscarriages or with a history of 1 or more term pregnancy followed by a minimum number of 2 consecutive miscarriages N = 54/54	- Medroxy Progesterone acetate - Oral - Not stated - 10 mg daily	Double blind, RCT vs placebo	Miscarriage, PTB
Klopper et al. 1965	Women who had 2 or more miscarriages, no pregnancy beyond 28 weeks' gestation, were less than 10 weeks into the current pregnancy and with no other obvious causes of miscarriage	- Cyclopentyl onol ether of Progesterone - Oral - Not stated - 50 mg twice daily	Double blind, RCT vs placebo	Miscarriage
Moller et al. 1965	Women with a positive pregnancy test. N = 40/40	- Medroxy Progesterone acetate - Oral - 20 mg/day for 3 days, followed by 10 mg/day for 11 days	Double blind, RCT vs placebo	Miscarriage

Moller et al. 1965	Women with a positive pregnancy test. N = 63/63	- Medroxy Progesterone acetate - Oral - 40 mg/day for 3 days, followed by 20 mg/day for 11 days	Double blind, RCT vs placebo	Miscarriage
Moller et al. 1965	Women with a positive pregnancy test. N = 153/153	- Medroxy Progesterone acetate - Oral - 80 mg/day for 3 days, followed by 40 mg/day for 11 days	Double blind, RCT vs placebo	Miscarriage
Levine et al. 1964	Women with 3 consecutive miscarriages of less than 16 weeks' gestation and with no signs of threatened miscarriage in the current pregnancy N = 56/30	- 17OHPc - IM - until miscarriage or 36 th week - 500 mg weekly	Double blind, RCT vs placebo	Miscarriage, PTB
Anderson et al. 2002	Women having undergone IVF or ICSI with a positive pregnancy test 14 days after transfer N = 303/303	- Progesterone vaginally - 3 weeks - 200 mg 3 times daily	RCT vs no treatment	Miscarriage
Reijnders et al. 1988	One or more of the following criteria: pregnancy after ovulation induction; 2 or more previous miscarriages; period of infertility for more than 12 months. Evidence of a viable fetus at 6 weeks of pregnancy N = 64/64	- 17OHPc - i.m. - from 7 th to 12 th week - 500 mg/week	Double blind, RCT vs placebo	Miscarriage, PTB
Shearman et al. 1963	Women with 2 or more consecutive abortions and low or falling pregnane-diol levels N = 50/50	- 17OHPc - i.m. - Up to 8 th week 250 mg/weekly; then increasing dose to 500 mg till 16 th week; then reducing doses to 250 mg/weekly until 24 th week	Double blind, RCT vs placebo	Miscarriage
Berle et al. 1980	Women up to 20 weeks gestation with bleeding N° = 300/300	- 90% received oral allylestrenol 15-20 mg/day - 10% received 250 mg of i.m. 17OHPc daily or every 2 days - Duration not stated	Double blind, RCT vs placebo	Miscarriage
Tognoni et al. 1980	Women with threatened miscarriage up to 14 weeks' gestation N° =145/139	- Oral allylestrenol 10 mg/d - 25 mg 17HP im every 5 d - 8 weeks	RCT vs placebo	Miscarriage
Gerhard et al. 1987	Women with vaginal bleeding in pregnancy and a closed cervical os N° = 64/35	- Progesterone - Vaginal suppositories - 25 mg twice/day	Double blind, RCT vs placebo	Miscarriage, birth weight, preterm labour
Swyer Daley 1953	Women with 2 or more consecutive miscarriages before 12 weeks' gestation N = 113/113	- 6 × 25 mg Progesterone pellets inserted in the gluteal muscle	RCT vs control group	Miscarriage, birth weight, PTB, stillbirth
Corrado et al. 2002	Women undergoing mid-second trimester amniocentesis N = 616/584	- Progesterone IM. 200 mg/day for 3 days followed by 17OHPc IM 340 mg twice/week - Duration unclear	RCT vs control group	Miscarriage, birth weight, preterm delivery

Table 2. — Characteristics of included studies for treating miscarriages.

Authors/year	Population studied	Intervention	Comparison	Outcomes measured
Gerhard et al. 1987	Women in the first trimester of pregnancy with confirmation of fetal viability by ultrasound scan N = 64/35.	Progesterone - Vaginal suppositories - 14 days after bleeding stopped - 25 mg twice/day	Double blind, RCT vs placebo	Miscarriage, birth-weight, preterm labour
Palagianio et al. 2004	50 women with previous diagnosis of inadequate luteal phase, threatened miscarriage and confirmed fetal viability	Progesterone - Vaginal suppositories - 5 days - 90 mg/day	RCT vs placebo	Pain relief uterine contractions, blood loss

Thirty-four studies were potentially eligible for inclusion. Only 2 were included after applying the inclusion criteria (Gerhard et al., 1987; Palagianio et al., 2004). The characteristics of these RCTs are described in Table 2.

These two trials met the inclusion criteria and involved 85 participants. Due to a paucity of data, subgroup analysis for early and late miscarriage, effect of progestagens by type, dose, and route of administration could not be carried out.

Discussion

The overall meta-analysis conducted on the use of progestagens for the treatment of recurrent miscarriages (Haas and Ramsey, 2009) showed no statistically significant difference in miscarriage rates between progestagens and placebo groups (Peto OR 0.98; 95% CI 0.78 -1.24). The analysis regarding the method of administration also showed no statistically significant difference between progestagens and placebo groups.

No evidence for an effect favouring the use of progestagens in women with recurrent miscarriage was found when compared to placebo (Peto OR 1.11, 95% CI 0.79-1.56, $P = 0.54$). Four studies (Shearman and Garrett, 1963; Levine, 1964; Reijnders et al., 1998; Corrado et al., 2002) compared intramuscular progestagens with placebo, the results showed no differences in miscarriages between the two groups (Peto OR 0.77; 95% CI 0.36-1.68, $P = 0.52$). Only one study compared vaginally administered progestagens with placebo, a second one comparing it with no treatment. The incidence of recurrent miscarriage was similar in both groups (Peto OR 0.74; 95% CI 0.40-1.35, $P = 0.32$).

The analysis concerning recurrent miscarriage considered 4 trials, of these 2 trials (Levine, 1964;

El-Zibdeh, 2005) enrolled only women who had suffered three or more miscarriages, 2 others (Swyer and Daley, 1953; Goldzieher, 1964) provided separate pregnancy outcome data by number of previous consecutive pregnancy losses. The results showed a reduction in miscarriage in favour of those in the progestagens group (Peto OR 0.38; 95% CI 0.20 to 0.70, $P = 0.002$). This finding however should be approached with caution as numbers are small.

Meta-analysis of the effect of vaginal Progesterone on miscarriage when compared to placebo (Wahabi et al., 2011) showed a point estimate which suggests a reduction of miscarriage rate with the use of Progesterone (risk ratio 0.47; 95% confidence interval 0.17 to 1.30), but the uncertainty about the outcome of this study is due to the small sample size.

Conclusion

Based on scarce data there is no evidence to support the routine use of progestagens for the treatment of threatened miscarriage. Information about potential harms to the mother or child, or both, with the use of progestagens is lacking. There has been much speculation about progestagens but the results of these meta-analysis show no statistically significant difference between women receiving progestagens for preventing threatened miscarriage or miscarriage when compared to placebo, other treatments or no treatment. Further, larger RCTs investigating potential harms as well as benefits, are urgently needed. A finding of a significantly reduced miscarriage rate in women with a history of recurrent miscarriage was found only in studies that included women with at least three miscarriages. This finding was obtained with different types of progestagens and therefore deserves further study before entering in clinical practice.

References

- Anderson NA, Popovic-Todorovic B, Schmidt K et al. Progesterone supplement during early gestations after IVF or ICSI has no effect on the delivery rates: a randomized controlled trial. *Hum Reprod* 2002;17:357-361.
- Berle P, Budenz M, Michaelis J. Is hormonal therapy still justified in imminent abortion? *Z Geburtshilfe Perinatol* 1980;184:353-358.
- Christian MS, Brent RL, Calda P. Embryo-fetal toxicity signals for 17alpha-hydroxyprogesterone caproate in high-risk pregnancies: a review of the non-clinical literature for embryo-fetal toxicity with progestins. *J Matern Fetal Neonatal Med.* 2007;20:89-112.
- Corrado F, Dugo C, Cannata M et al. A randomised trial of progesterone prophylaxis after midtrimester amniocentesis. *Eur J Obstet Gynecol Reprod Biol* 2002;100:196-198.
- Cunningham FG. Reproductive success and failure. *Williams obstetrics*. Section IX. 21st Edition. New York: McGraw-Hill, 2001.
- Di Renzo GC, Mattei A, Gojnic M et al. Progesterone and Pregnancy. *Current opinion Obstet Gynecol.* 2005;17:598-600.
- El-Zibdeh MY. Dydrogesterone in the reduction of recurrent spontaneous abortion. *J Steroid Biochem Mol Biol* 2005; 97:431-434.
- Everett C. Incident and outcome of bleeding before the 20th week of pregnancy: prospective study from general practice. *BMJ* 1997;315:32-34.
- Gerhard I, Gwinner B, Eggert-Kruse W et al. Double-blind controlled trial of progesterone substitution in threatened abortion. *Biol Res in Pregnancy Perinatol* 1987;8:26-34.
- Goldzieher JW. Double-blind trial of a progestin in habitual abortion. *JAMA* 1964;188:651-654.
- Haas DM, Ramsey PS. Progestogen for preventing miscarriage. *Cochrane Database Syst Rev.* 2009; CD003511.
- Hemminki E. Treatment of miscarriage: current practice and rationale. *Obstet Gynecol* 1998;91:247-253.
- Klopper A, MacNaughton. M. Hormones in recurrent abortion. *J Obstet Gynaecol Br Commonw* 1965;72:1022-1028.
- Levine L. Habitual abortion. A controlled clinical study of progestational therapy. *West J Surg Obstet Gynecol* 1964; 72:30-36.
- Moller K, Fuchs F. Double blind controlled trial of 6- methyl-17-acetoxypregesterone in threatened abortion. *J Obstet Gynaecol Br Commonw*1965;72:1042-1044.
- Mosby Incorporated. Medroxyprogesterone acetate. Harcourt Health Sciences Incorporated, 2001.
- Palagiano A, Bulletti C, Pace MC et al. Effects of vaginal progesterone on pain and uterine contractility in patients with threatened abortion before twelve weeks of pregnancy. *Ann N Y Acad Sci* 2004;1034:200-210.
- Reijnders FJL, Thomas CMG, Doesburg WH et al. Endocrine effects of 17 alphahydroxyprogesterone caproate during early pregnancy: a double-blind clinical trial. *Br J Obstet Gynaecol* 1988;95:462-468.
- Shearman R, Garrett W. Double-blind study of effect of 17-hydroxyprogesterone caproate on abortion rate. *BMJ* 1963;1:292-295.
- Swyer GIM, Daley D. Progesterone implantation in habitual abortion. *BMJ* 1953;1:1073-1086.
- Szabo I, Szilagy A. Management of threatened abortion. *Early Pregnancy* 1996;2:233-240.
- Tognoni G, Ferrario L, Inzalaco M et al. Progestogens in threatened abortion. *Lancet* 1980;2: 1242-1243.
- Wahabi HA, Abed Althagafi NF, Elawad M, Al Zeidan RA. Progestogen for treating threatened miscarriage. *Cochrane Database Syst Rev* 2011;(3):CD005943.
- Whitehead MI, Hillard TC, Crook D. The role and use of progestogens. *Obstet Gynecol* 1990;75(4 Suppl):59S-76S; discussion 81S-83S.
- World Health Organization. *International Statistical Classification of Diseases and Related Health Problems*, 10th Edition.,Vol. 1, Geneva, 1992.